

Finite-size energy levels of the superintegrable chiral Potts model

G. von Gehlen

Physikalisches Institut der Universität Bonn, Nussallee 12, 53115 Bonn, Germany

Abstract. In the solution of the superintegrable chiral Potts model special polynomials related to the representation theory of the Onsager algebra play a central role. We derive approximate analytic formulae for the zeros of particular polynomials which determine sets of low-lying energy eigenvalues of the chiral Potts quantum chain. These formulae allow the analytic calculation of the leading finite-size corrections to the energy eigenvalues without resorting to a numerical determination of the zeros.

1 Introduction

The chiral Z_N -symmetrical Potts model, apart from showing a rich phase structure, has attracted considerable interest because for appropriate parameter choices it has very special integrability properties. In the "superintegrable" case it is integrable *both* because it provides a representation of Onsager's algebra (in this respect being a natural generalization of the Ising model), and also because its Boltzmann weights satisfy a new type of Yang-Baxter equation. Both properties guarantee the existence of an infinite set of commuting charges. In order to calculate the spectrum, functional relations are needed. Such relations had first been conjectured by Albertini *et al.* (1989), and later derived by exploring relations to the six-vertex model at $q = e^{2\pi i/N}$ (Bazhanov and Stroganov 1990, Baxter *et al.* 1990). The formulae which give the spectrum in the superintegrable case have been given by Albertini *et al.* (1989) for Z_3 and for general Z_N by Baxter (1994). Little is known about correlation functions. For the general integrable case (not satisfying the Onsager algebra), the free energy and interface tensions have been obtained in the thermodynamic limit (O'Rourke and Baxter 1996). It is still a challenge to find an analytic derivation of the order parameter (Baxter 1998). Recently, general integrable boundary conditions for the chiral Potts model have been obtained by Zhou (1998).

In the following we concentrate on the solution of the superintegrable case, in which special polynomials related to the representation theory of the Onsager algebra play a central role. The energy eigenvalues are determined in terms of the zeros of these polynomials. For the lowest states the polynomials are of order $m_E = [((N-1)L - Q)/N]$. L denotes the number of sites of the chain and Q the Z_N -charge sector. $[x]$ stands for the integer part of a rational number x . In the limit $L \rightarrow \infty$ the sum over the terms depending on the zeros can be expressed as a contour integral without explicitly calculating the zeros. This method has been used to obtain the ground state energy and excited levels

in the thermodynamic limit. For the gapless regime finite-size corrections have been studied by Albertini and McCoy (1990) in order to obtain information on correlation functions through arguments of conformal invariance. Interesting effects due to the broken parity invariance have emerged. For finite L one has to resort to numerical methods.

In these lecture notes, after reviewing main features of the model and its solution, we study the zeros of the above mentioned polynomials. We derive quite accurate approximate analytic expressions for their location at finite L .

2 Integrability of the chiral Potts model

2.1 The integrable model

For many discussions of the chiral Potts quantum chain, it turns out to be sufficient to consider the following Z_N -symmetric hamiltonians which depend on three real parameters λ , ϕ , φ :

$$H = - \sum_{j=1}^L \sum_{l=1}^{N-1} \frac{1}{\sin(\pi l/N)} \left(e^{i\varphi(2l-N)/N} Z_j^l + \lambda e^{i\phi(2l-N)/N} X_j^l X_{j+1}^{N-l} \right).$$

The first sum is over the sites j . At each site j there are operators X_j and Z_j acting in a vector space C^N and satisfying

$$Z_i X_j = X_j Z_i \omega^{\delta_{i,j}}; \quad Z_j^N = X_j^N = 1; \quad \omega = e^{2\pi i/N}.$$

Often it is useful to represent the operators X_j and Z_j by $(X_j)_{l,m} = \delta_{l,m+1} \pmod{N}$ and $Z_j = \text{diag}(1, \omega, \omega^2, \dots, \omega^{N-1})_j$. H commutes with the Z_N -charge operator $\hat{Q} = \prod_{j=1}^L Z_j$. We write the eigenvalues of $\hat{Q}$ as ω^Q where $Q = 0, 1, \dots, N-1$. λ is the inverse temperature. We shall consider the periodic case $X_{N+1} = X_1$, although, as shown by Baxter (1994), twisted toroidal boundary conditions require little additional effort.

2.2 Particular cases: Ising, Parafermionic

For $N = 2$ the sum over l has only a single term. The angles ϕ , φ drop out, Z_j and X_j become Pauli matrices and we obtain the Ising quantum chain. If we put $\phi = \varphi = 0$ then H describes the Fateev-Zamolodchikov parity invariant parafermionic Z_N -quantum chain. For $N \geq 3$ and $\phi \neq 0$ or $\varphi \neq 0$ parity is broken (therefore ϕ and φ are called chiral angles) and this gives rise to several interesting features of the model, e.g. to the appearance of incommensurate phases and oscillating correlation functions.

2.3 Yang-Baxter-integrable case

If $\cos \varphi = \lambda \cos \phi$ then H can be derived from a two-dimensional lattice model defined by the following transfer matrix which depends on rapidities p and q :

$$T_{p,q}(\{l\}, \{l'\}) = \prod_{j=1}^L W_{p,q}(l_j - l'_j) \bar{W}_{p,q}(l_j - l'_{j+1}) \quad (1)$$

$\{l\}$ and $\{l'\}$ are the sets of N -valued spin variables at alternating vertex rows of the diagonally drawn lattice. The rapidity lines p run horizontally on the dual lattice, the q vertically. The Boltzmann weights W and $\bar{W}$ are defined in terms of the functions $x_p, y_p, \mu_p; x_q, y_q, \mu_q$ of the rapidities p and q by (Au-Yang *et al.* 1987)

$$W_{p,q}(n) = \left(\frac{\mu_p}{\mu_q} \right)^n \prod_{j=1}^n \frac{y_q - \omega^j x_p}{y_p - \omega^j x_q}; \quad \bar{W}_{p,q}(n) = (\mu_p \mu_q)^n \prod_{j=1}^n \frac{\omega x_p - \omega^j x_q}{y_q - \omega^j y_p}$$

where $n = 0, 1, \dots, N-1$. The requirement of Z_N -symmetry imposes several restrictions on the six functions, e.g.

$$\lambda(x_q^N y_q^N + 1) = x_q^N + y_q^N.$$

where λ is a parameter describing the inverse temperature. At fixed λ only one of the three functions x_q, y_q, μ_q is independent, the other two are determined by nonlinear relations.

In the Ising case $N = 2$, the functions x_q etc. can be simply expressed in terms of the meromorphic elliptic functions of modulus λ :

$$x_q = -\sqrt{\lambda} \operatorname{sn} q; \quad y_q = \sqrt{\lambda} \operatorname{cn} q / \operatorname{dn} q, \quad \mu_q = \sqrt{1 - \lambda^2} / \operatorname{dn} q$$

and one obtains the parametrization of the Ising Boltzmann weights known e.g. from Baxter (1982). Integrability is guaranteed by the following Yang-Baxter-equation, which contains the p - and q -variables *separately*

$$\begin{aligned} \sum_{l=0}^{N-1} W_{p,q}(l' - l) \bar{W}_{r,q}(l'' - l) W_{r,p}(l - l''') \\ = R_{pqr} \bar{W}_{r,p}(l'' - l') W_{r,q}(l' - l''') \bar{W}_{p,q}(l'' - l'''). \end{aligned}$$

The explicit expression for R_{pqr} can be found e.g. in Albertini *et al.* (1989). The superintegrable case corresponds to fixing the p -dependent functions to $x_p = y_p$ and $\mu_p = 1$ so that we are left with only one independent function, e.g. x_q . H is obtained if at fixed p we expand the transfer matrix for small $q - p$.

2.4 The superintegrable case, Onsager's algebra

The choice $\phi = \varphi = \pi/2$ gives the "superintegrable" case of the hamiltonian. Writing $H = \frac{1}{2}N(A_0 + \lambda A_1)$ we have

$$A_0 = -\frac{4}{N} \sum_{j=1}^L \sum_{l=1}^{N-1} \frac{Z_j^l}{1 - \omega^{-l}}; \quad A_1 = -\frac{4}{N} \sum_{j=1}^L \sum_{l=1}^{N-1} \frac{X_j^l X_{j+1}^{N-l}}{1 - \omega^{-l}}. \quad (2)$$

It has been shown by v.Gehlen and Rittenberg (1985) that A_0 and A_1 satisfy the Dolan and Grady (1982) conditions

$$[A_0, [A_0, [A_0, A_1]]] = 16 [A_0, A_1]; \quad [A_1, [A_1, [A_1, A_0]]] = 16 [A_1, A_0]. \quad (3)$$

Due to these conditions, A_0 and A_1 generate Onsager's algebra $\mathcal{A}$:

$$[A_l, A_m] = 4 G_{l-m}$$

$$[G_l, A_m] = 2 A_{m+l} - 2 A_{m-l}; \quad [G_l, G_m] = 0; \quad l, m \in \mathbb{Z}.$$

which has been an essential tool in the Onsager (1944) original solution of the Ising model. $\mathcal{A}$ implies an infinite set of constraints:

$$[A_2, A_1] = [A_1, A_0] = [A_0, A_{-1}] = \dots$$

and the existence of the infinite set of commuting charges Q_m :

$$Q_m = \frac{1}{2} (A_m + A_{-m} + \lambda(A_{m+1} + A_{-m+1})) \quad \text{with} \quad Q_0 = H.$$

Finite dimensional representations of $\mathcal{A}$ are obtained if there is a recurrence relation of finite length among the A_l (Davies 1990, Roan 1991):

$$\sum_{k=-n}^n \alpha_k A_{k-n} = 0.$$

If z_j ($j = 1, \dots, m_E$) are the zeros of the polynomial $f(z) = \sum_{k=-n}^n \alpha_k z^{k+n}$, then we can express the A_m and G_m in terms of a set of $sl(2, C)$ -generators

$$A_m = 2 \sum_{j=1}^n (z_j^m E_j^+ + z_j^{-m} E_j^-); \quad G_m = \sum_{j=1}^n (z_j^m - z_j^{-m}) H_j$$

where

$$[E_j^+, E_k^-] = \delta_{jk} H_k; \quad [H_j, E_k^\pm] = \pm 2 \delta_{jk} E_k^\pm.$$

So, $\mathcal{A}$ is isomorphic to a subalgebra of the loop algebra of a direct sum of $sl(2, C)$ algebras.

Each sector of the model is characterized by a set of points z_j , which, of course, are not fixed by the algebra $\mathcal{A}$ alone. For the hamiltonian (2) the polynomials $f(z)$, or rather related polynomials $\mathcal{P}_Q(s)$, have been obtained by Albertini *et al.* (1989) and Baxter (1994) from functional equations for the transfer matrix

(1) of the two-dimensional model, as will be reviewed in the next section. As B. Davies (1990) has shown, from the property (3) of H it follows that *all* eigenvalues E of H depend on λ in the form

$$E = a + b\lambda + N \sum_{j=1}^{m_E} \pm \sqrt{1 + 2\lambda \cos \theta_j + \lambda^2} \quad (4)$$

where a and b are integers and $\cos \theta_j = \frac{1}{2} (z_j + z_j^{-1})$. So the main task of the diagonalization of the superintegrable H reduces to the calculation of the $\cos \theta_j$ or equivalently, of the zeros of the polynomials $f(z)$. For hermitian hamiltonians, the z_j must be on the unit circle.

3 Solution of the Z_3 -superintegrable model

3.1 Functional equation and ansatz of Albertini et al.

Albertini *et al.* (1989) discovered the following functional equation for the Z_3 superintegrable case with the p -variables fixed to $x_p = y_p = \left(\frac{1-\lambda}{1+\lambda}\right)^{1/6}$: Splitting off some poles from $T_{p,q}$ they get a meromorphic function $\bar{T}_q$ which satisfies

$$\begin{aligned} \bar{T}_q \bar{T}_{Rq} \bar{T}_{R^2q} = & 3^L e^{-iP} \{ (t-1)^L (\omega^2 t - 1)^L \bar{T}_q \\ & + (\omega t - 1)^L (\omega^2 t - 1)^L \bar{T}_{R^2q} + (t-1)^L (\omega t - 1)^L \bar{T}_{R^4q} \} \end{aligned} \quad (5)$$

where P is the momentum operator, $t = x_q y_q / x_p^2$ and R is the mapping $R(x_q, y_q, \mu_q) = (y_q, \omega x_q, \mu_q^{-1})$. Solutions to this functional equation are then obtained from the ansatz

$$\bar{T}_q = \dots \prod_{l=1}^{m_P} \left(\frac{1 + \omega v_l t}{1 + \omega v_l} \right) \prod_{j=1}^{m_E} \{ (1 - \lambda)(\mu_q^3 + 1) \pm 2w_j(\mu_q^3 - 1) \}. \quad (6)$$

We skip here writing some factors relevant for the determination of the linear terms in (4) and chose a certain integer $P_b = 0$: for details see Albertini and McCoy (1990) or Baxter (1993,1994). The ansatz (6) contains m_P excitations with Bethe-type rapidities v_l , and in the last factor, m_E functions w_j . It requires some calculation to see that if we put

$$w_j = \frac{1}{2} \sqrt{1 + 2\lambda \frac{1 + t_j^3}{1 - t_j^3} + \lambda^2}$$

then the left hand side of (5) vanishes, and in order to get the right hand side vanishing too, t_j must be zero of the polynomial

$$\begin{aligned} \mathcal{P}_Q(t^3) = & \frac{t^{-c}}{3} \left\{ (t^2 + t + 1)^L \prod_{l=1}^{m_P} \frac{1 + v_l t}{1 + v_l^3 t^3} + \omega^Q (t^2 + \omega^2 t + \omega)^L \prod_{l=1}^{m_P} \frac{1 + \omega v_l t}{1 + v_l^3 t^3} \right. \\ & \left. + \omega^{2Q} (t^2 + \omega t + \omega^2)^L \prod_{l=1}^{m_P} \frac{1 + \omega^2 v_l t}{1 + v_l^3 t^3} \right\} \end{aligned} \quad (7)$$

The integer c is chosen such that $\mathcal{P}_Q$ becomes invariant against $t \rightarrow \omega t$ and $\mathcal{P}_Q$ depends only on $s \equiv t^3$ with $\mathcal{P}_Q(0)$ finite. Despite its denominators, $\mathcal{P}_Q(s)$ will be a polynomial if the v_l satisfy the Bethe-type equations:

$$\left(\frac{1 + \omega v_j}{1 + \omega^2 v_j} \right)^L = (-1)^{m_P+1} \omega^{Q-m_P} \prod_{l=1}^{m_P} \left(\frac{v_l - \omega v_j}{v_j - \omega v_l} \right).$$

These Bethe-type equations resemble those of the spin-1 XXZ-chain. As mentioned already above, the hamiltonian is obtained by expanding the transfer matrix for small $p - q$ around a fixed value of p . If $\mathcal{P}_Q$ has degree m_E , then from the m_E zeros of (7) we get a set of 2^{m_E} energy eigenvalues

$$E_i = a(1 + \lambda) + b - 6(\pm w_1 \pm w_2 \pm \dots \pm w_{m_E}). \quad (8)$$

where e.g. for the lowest sector $a = 3m_E - 2L$, $b = 2Q$. Comparing to the formula eq.(4) obtained from the Onsager algebra, we see that $\cos \theta_j = (1 + t_j^3)/(1 - t_j^3)$. Our hamiltonian is hermitian and so we should have $|\cos \theta_j| \leq 1$ and all zeros $s_j = t_j^3$ must be on the negative real s -axis.

A difficult question is whether all solutions to these equations really give eigenvalues of our H , or whether there are spurious solutions. This problem has been investigated in detail by Dasmahapatra *et al.* (1993).

3.2 Example: Sector $m_P = 0$

In general, in order to obtain energy eigenvalues of H for chain length L in the charge sector Q , one first has to choose a sector of given m_P and to solve the Bethe-type equations obtaining a set of Bethe-rapidities v_l . Then one has to use these v_l to build up $\mathcal{P}_Q$ and then to calculate its zeros s_j .

Obviously, the simplest case is the sector $m_P = 0$ where there are no v_l and one can directly start with building the $\mathcal{P}_Q$ and looking for the zeros s_j . E.g. for Z_3 with $L = 20$ and $Q = 1$ we calculate first

$$\begin{aligned} \mathcal{P}_1(s) = & 20 s^{13} + 8455 s^{12} + 484500 s^{11} + 8533660 s^{10} + 61757600 s^9 \\ & + 210859245 s^8 + 363985680 s^7 + 326527350 s^6 + 151419816 s^5 \\ & + 34880770 s^4 + 3656360 s^3 + 146490 s^2 + 1520 s + 1, \end{aligned}$$

then solve numerically for its 13 zeros. Since the largest appearing coefficients grow exponentially with L (all coefficients are positive, and from (7) with $m_P = 0$ we get $\mathcal{P}_Q(1) = 3^{L-1}$), this procedure becomes impractical for $L \lesssim 50$. and e.g. energy eigenvalues for $L = 1000$ can hardly be obtained this way. Can we get an analytic expression for the zeros of the $\mathcal{P}(s)$ for large *finite* L ?

Baxter (1988) and Albertini *et al.* (1989) have obtained closed expressions for energy levels *in the limit* $L \rightarrow \infty$. One method to get these is to rewrite the sum over the terms determined by the zeros (choosing all $\pm$ -signs to be + and so

obtaining the lowest eigenvalue E) as an integral over a contour C enclosing the negative real t -axis:

$$\sum_{j=1}^{m_E} \sqrt{1 + 2\lambda \cos \theta_j + \lambda^2} = -\frac{3}{2\pi i} \int_C dt \sqrt{(1 - \lambda)^2 + \frac{4\lambda}{1 - t^3}} \frac{d}{dt} \ln \mathcal{P}_Q(t)$$

Then the contour is opened and gets caught by three 120° symmetrically oriented cuts, which arise from the square root of the integrand. The rightmost of these cuts is on the positive real axis at $1 \leq t \leq |(1 + \lambda)/(1 - \lambda)|^{1/3}$. All three cuts give the same contribution, and the rightmost one can be evaluated easily, since for $L \rightarrow \infty$ only the term proportional $(t^2 + t + 1)^L$ of $\mathcal{P}$, eq.(7), is non-oscillating and needs to be kept. Baxter (1988) finds that the ground-state energy for general Z_N is given by (see also Albertini *et al.* 1989):

$$\lim_{L \rightarrow \infty} \frac{E_0}{L} = -\frac{N}{\pi} \int_0^{\pi(1-N^{-1})} dx \sqrt{1 - 2\lambda \frac{\sin^N(x + \frac{\pi}{N}) - \sin^N x}{\sin^N(x + \frac{\pi}{N}) + \sin^N x}} + \lambda^2. \quad (9)$$

4 Determination of the zeros of the $\mathcal{P}_Q(s)$

We attempt to find the zeros of the $\mathcal{P}_Q(s)$ for *finite* L on the negative real s -axis analytically. From the explicit expressions and numerical studies we observe:

- For $L = Q \bmod 3$ the zeros come in pairs which are reflection symmetric with respect to $s = -1$: with s_j also $1/s_j$ is a zero. This remains approximately (within a shift) true for the other charge sectors.
- The distances between adjacent zeros alternate slightly in size so that a smooth curve $s_j(k)$ is obtained only if we connect every second zero.

4.1 Case Z_3 , $m_P = 0$

A useful variable transformation to be applied to $\mathcal{P}_Q(s)$ turns out to be $t = (u - \sqrt{3})/(u + \sqrt{3})$ or, writing $u = \tan \beta$:

$$t = \frac{\sin(\beta - \frac{\pi}{3})}{\sin(\beta + \frac{\pi}{3})}.$$

This maps $-\infty < s < 0$ to e.g. $-\frac{\pi}{3} < \beta < \frac{\pi}{3}$ or $\frac{2\pi}{3} < \beta < \frac{4\pi}{3}$ (for definiteness let us choose the first interval), giving

$$t^2 + t + 1 = \frac{3}{(2 \sin(\beta + \frac{\pi}{3}))^2} \quad \text{and} \quad t^2 + \omega t + \omega^2 = \frac{-3 \omega \cos \beta e^{-i\beta}}{2 (\sin(\beta + \frac{\pi}{3}))^2}.$$

so that

$$\mathcal{P}_Q \sim (t^2 + t + 1)^L + \omega^Q (t^2 + \omega^2 t + \omega)^L + \omega^{-Q} (t^2 + \omega t + \omega^2)^L = 0,$$

multiplying by the non-vanishing factors $(\sin(\beta + \frac{\pi}{3}))^{2L}$ becomes




Fig. 1. Left-hand (dash-dotted line) and right hand (dotted and dashed) parts of (10) for $L = 12$ sites. Charge sector $Q = 0$: dotted lines, sector $Q = 1$: dashed lines. The intersections with the dash-dotted line with $-\pi/3 < \beta < \pi/3$ give the m_E solutions β_k : $m_E = 8$ for $Q = 0$ and $m_E = 7$ for $Q = 1$.

$$-(2 \cos \beta)^{-L} = 2 \cos \left(L \left\{ \beta + \frac{\pi}{3} \right\} + \frac{2\pi Q}{3} \right). \quad (10)$$

Now $-\frac{\pi}{3} < \beta < \frac{\pi}{3}$ means $2 \cos \beta > 1$ so that for $L \gg 1$ the left hand side of (10) will be small, except near the corners of the interval in β , as is shown in Fig. 1. In order to get a solution, the fast oscillating right hand term has to coincide with the non oscillating left hand term. We see that all but the solutions near $\beta_k \approx -\frac{\pi}{3}$ and $\beta_k \approx \frac{\pi}{3}$ are practically equally spaced in the variable β . Neglecting the left-hand side altogether, we get the approximate solutions

$$\beta_k \approx K - \frac{\pi}{3}; \quad K = \frac{6k + 2Q - 3}{6L} \pi; \quad k = 1, 2, \dots, m_E. \quad (11)$$

or, expressed in terms of the $\cos \theta_k$ which we need for the energy eigenvalues:

$$\cos \theta_k \equiv \frac{1 + t_k^3}{1 - t_k^3} = -\frac{\sin^3(K + \frac{\pi}{3}) - \sin^3 K}{\sin^3(K + \frac{\pi}{3}) + \sin^3 K}. \quad (12)$$

It is easy to see that using (11, 12) and replacing the summation in (8) by an integral leads to Baxter's formula (9) for the thermodynamic limit. Through a Q -dependent shift (11) describes most of the Q -dependence of the finite- L levels. Further finite-size effects are due to the deviations of the largest and smallest β_k from the approximation (11).

In order get an idea of the precision of (11), consider e.g. $L = 30$, $Q = 0$. With $d_k = (\beta_k^{appr} / \beta_k^{exact}) - 1$ where β_k^{appr} is the approximation (11), we find

$$d_1 = 0.0012; d_2 = -1.89 \cdot 10^{-5}; d_3 = 6.48 \cdot 10^{-7}; d_4 = -4.39 \cdot 10^{-8}; \dots$$

For $L = 30000$ and $Q = 0$ we get $d_1 = 0.99 \cdot 10^{-6}$, but compared to $L = 30$ also the spacing between neighbouring β_k is $\sim 2 \cdot 1000/3$ times smaller.

Taking the arcsin of (10) we get a pair of useful (still exact) equations for the β_k :

$$\mp \arcsin \frac{1}{2(2 \cos \beta_k)^L} + L \left(\beta_k + \frac{\pi}{3} \right) + \frac{2\pi Q}{3} - 2\pi I_k^\pm \mp \frac{\pi}{2} = 0. \quad (13)$$

The two sets of counting integers $I_k^\pm$ vary within the limits

$$I_k^\pm = I_a^\pm, I_a^\pm + 1, \dots, I_b^\pm$$

with

$$I_a^\pm = \left\lfloor \frac{Q + 3 \mp 1}{3} \right\rfloor; \quad I_b^\pm = \left\lfloor \frac{L + Q - 1 \mp 1}{3} \right\rfloor; \quad (14)$$

The arcsin-equation has the great advantage that from it each zero can be calculated separately by fixing $I_k^\pm$. Numerically on a small PC we can easily get zeros with 40 digits precision for up to $L = 300000$ sites. Recall that computing directly the zeros of the $\mathcal{P}_Q(t)$ we were limited to $L \lesssim 50$. In addition, the arcsin-equation lends itself to excellent analytical approximations for the "corner"-solutions: Let us call ξ the term which is important only close to the corners:

$$\xi \equiv \arcsin \frac{1}{2(2 \cos \beta)^L} \quad (15)$$

We find numerically that $\xi_k^\pm \ll 1$ even for the extreme cases $I_k^\pm = I_{a,b}^\pm$. We use this feature to approximate ξ_k . Let us consider the lower end of the interval in β , i.e. $I_k \gtrsim I_a$. Then we get a behaviour smooth in L when keeping Q fixed (At the upper end $I_k \lesssim I_b$ the behaviour is smooth only if we keep $L + Q$ fixed mod 3). Using $\arcsin x \approx x$ and writing $\beta = -\pi/3 + \epsilon$ we get (omitting the superscripts $\pm$):

$$\xi \approx \frac{1}{2} (2 \cos \beta)^{-L} = \frac{1}{2} \left(\cos \epsilon + \sqrt{3} \sin \epsilon \right)^{-L} \approx \frac{1}{2} \left(1 + \sqrt{3} \epsilon \right)^{-L}. \quad (16)$$

On the other hand, solving (13) for the β_k which is not in ξ_k , we have:

$$\epsilon_k = \frac{\Gamma_k \pm \xi_k}{L} \quad \text{with} \quad \Gamma_k = \left(2I_k - \frac{2Q}{3} \pm \frac{1}{2}\right) \pi. \quad (17)$$

Inserting (17) into (16) and assuming $L \gg 1$ we get

$$\xi_k \approx \frac{1}{2} \left(1 + \frac{\sqrt{3}(\Gamma_k \pm \xi_k)}{L}\right)^{-L} \approx \frac{1}{2} \exp\left\{-\sqrt{3}(\Gamma_k \pm \xi_k)\right\} \quad (18)$$

This nonlinear equation for ξ_k , valid for small ϵ , can be solved by iteration. The zeroth and first iterations give

$$\xi_k^{\pm(0)} = \frac{1}{2} e^{-\sqrt{3}\Gamma_k^{\pm}} \quad \text{and} \quad \xi_k^{\pm(1)} = \frac{1}{2} \exp\left\{-\sqrt{3}\left(\Gamma_k^{\pm} \pm \frac{1}{2} e^{-\sqrt{3}\Gamma_k^{\pm}}\right)\right\},$$

respectively, or

$$\beta_k^{\pm(1)} = -\frac{\pi}{3} + \frac{\Gamma_k^{\pm}}{L} \pm \frac{1}{2L} \exp\left\{-\sqrt{3}\left(\Gamma_k^{\pm} \pm \frac{1}{2} e^{-\sqrt{3}\Gamma_k^{\pm}}\right)\right\}, \quad \text{etc.}$$

So we find corrections to (11) which fast become smaller with increasing I_k , and are useful mainly for two smallest values β_k . Closer to the middle of the interval, anyway the approximation $\xi_k = 0$ gets very accurate.

Coming back to the example $L = 30000$, $Q = 0$ given above, adding the i th iteration corrections $\xi_k^{(i)}/L$ improves the approximation (11) to

$$d_1^{(0)} = -5.48 \cdot 10^{-8}, \quad d_1^{(1)} = 3.285 \cdot 10^{-9}, \quad d_1^{(2)} = 1.53 \cdot 10^{-10}$$

for the zeroth, first and second iteration, respectively. Regarding the β_k away from the corners, already from (11) we get e.g. $d_7 = 1.4 \cdot 10^{-25}$, which improves slightly to $d_7^{(0)} = 6.3 \cdot 10^{-26}$ by adding the zeroth iteration.

Let us study how much the results for the ξ_k are affected by the approximations made in (16) and (18), and how many iterations are needed. In Table 1 we give numerical values from our iteration formula in order to compare them to the following numerical results from the exact formula (13). A fit of the L -dependence of the numerical solution, obtained from $L = 25000, \dots, 125000$ sites is, expressed in terms $\xi^{\pm}/\pi$:

$$\begin{aligned} Q = 0+ : & +0.009927623657298 + 0.04835895/L + 0.0315/L^2 + \dots \\ Q = 0- : & -0.000045412321296 - 0.002017284/L - 0.0338/L^2 - \dots \\ Q = 1+ : & +0.00000740160385053 + 0.0004910486/L + 0.0130/L^2 + \dots \\ Q = 1- : & -0.0017242062241336 - 0.023760499/L - 0.0922/L^2 - \dots \\ Q = 2+ : & +0.0002780560828110 + 0.007462876/L + 0.0685/L^2 + \dots \\ Q = 2- : & -0.00000120676616449 - 0.000111825/L - 0.00430/L^2 - \dots \end{aligned}$$

We see that for large L our equation (18) determines the correction (15) to (11) with excellent precision if its twice iterated solution is used. The coefficients in

these expansions indicate that the asymptotic large- L -behaviour sets in only for about $L \gtrsim 1000$.

If we move downwards from the upper corner $\beta = \pi/3$, we get exactly the same numbers and formulae, just signs are changed and charge sectors are permuted.

Table 1. Values of ξ_k for the lowest value of β_k , as obtained from (18) at different iteration levels. We have multiplied the values by $10^6/\pi$ in order to avoid too small numbers.

Iteration	$Q = 0 +$	$Q = 0 -$	$Q = 1 +$	$Q = 1 -$	$Q = 2 +$	$Q = 2 -$
0st	10477.0	-45.401101	7.401901956	-1708.097	278.47707	-1.20675824029
1st	9896.39	-45.412318	7.401603838	-1724.047	278.05541	-1.20676616444
2nd	9927.70	-45.412321	7.401603850	-1724.196	278.05605	-1.20676616449

4.2 The leading finite-size corrections

The formulae of the last section are sufficient to calculate finite-size eigenvalues. But often it is convenient to transform the sums in (8) into integrals in order to have a compact expression also for finite L eigenvalues or to get the leading finite-size corrections to the thermodynamic limit results. This can be done e.g. applying Euler-Maclaurin-techniques as is common in calculations for the XXZ-quantum chain. Following Woynarovich and Eckle (1987) and Hamer *et al.* (1987), but taking care of the alternating spacing of the zeros, we define two functions $Z_L^\pm$

$$Z_L^\pm(\beta) = \frac{v_L^\pm}{2\pi L} \left\{ \mp \arcsin \frac{1}{2(2 \cos \beta)^L} + L \left(\beta + \frac{\pi}{3} \right) + \frac{2\pi Q}{3} \mp \frac{\pi}{2} \right\},$$

where $v_L^\pm = 3m_E^\pm/L$ so that $\lim_{L \rightarrow \infty} v_L^\pm = 1$. From (14) we read off $m_E^\pm = I_b^\pm - I_a^\pm + 1$. In $Z_L^\pm$ the roots β_k are exactly equidistant. We define the root density

$$\sigma_L^\pm(\beta) = dZ_L^\pm/d\beta = \frac{v_L^\pm}{2\pi} \left\{ 1 \mp (-1)^L \frac{\tan \beta}{\sqrt{4(2 \cos \beta)^{2L} - 1}} \right\},$$

and use $\sigma_\infty^+(\beta) + \sigma_\infty^-(\beta) = \sigma_\infty$, $m_E^+ + m_E^- = m_E$. Then the finite-size correction to a quantity $\int d\beta \sigma_\infty w(\beta)$ (since we are mainly interested in the ground-state energy per site, this will be the w of (8)) is

$$\frac{1}{L} \left(\sum_{k=1}^{m_E^+} w(\beta_k^+) + \sum_{k=1}^{m_E^-} w(\beta_k^-) \right) - \int_{-\pi/3}^{\pi/3} d\beta \sigma_\infty w(\beta)$$

$$\begin{aligned}
&= \int_{-\pi/3}^{\pi/3} d\beta (\sigma_L^+ + \sigma_L^- - \sigma_\infty) w(\beta) - \sum_{\pm} \left(\int_{-\pi/3}^{\beta_a^\pm} + \int_{\beta_b^\pm}^{\pi/3} \right) d\beta \sigma_L^\pm w(\beta) \\
&+ \frac{1}{2L} \sum_{\pm} \{w(\beta_b^\pm) + w(\beta_a^\pm)\} + \frac{1}{12L^2} \sum_{\pm} \left(\frac{w'(\beta_b^\pm)}{\sigma_L^\pm(\beta_b^\pm)} - \frac{w'(\beta_a^\pm)}{\sigma_L^\pm(\beta_a^\pm)} \right) + \dots \quad (19)
\end{aligned}$$

Both $\beta_a^\pm \equiv \beta_1^\pm = \beta^\pm(I_a^\pm)$ and $\beta_b^\pm = \beta^\pm(I_b^\pm)$ depend on the charge Q .

5 The zeros for higher Z_N -superintegrable cases

Analogous trigonometric formulae result for the higher Z_N -cases. With increasing N , the equations get more involved, but have a similar structure.

5.1 The case Z_4 :

Since we transform the variable t and not $s = t^4$, in order to map the negative real s -axis to a range of real β , we have to include a phase and use

$$t = e^{i\pi/4} \frac{\sin(\beta - \frac{\pi}{4})}{\sin(\beta + \frac{\pi}{4})} \quad (20)$$

so that $-\infty < s < 0$ is mapped to e.g. $-\frac{\pi}{4} < \beta < \frac{\pi}{4}$. After some algebra, we get that the condition for the zeros of the $m_P = 0$ -functions $\mathcal{P}_Q(s)$

$$\begin{aligned}
\mathcal{P}_Q(s) = t^{-c} \{ &(t^3 + t^2 + t + 1)^L + (-1)^Q (t^3 - t^2 + t - 1)^L \\
&+ i^{-Q} (t^3 + it^2 - t - i)^L + i^Q (t^3 - it^2 - t + i)^L \} = 0 \quad (21)
\end{aligned}$$

translates to

$$\begin{aligned}
&\left(\frac{\sqrt{2} - \cos(2\beta)}{\sqrt{2} + \cos(2\beta)} \right)^{L/2} \cos \left\{ L \left(\rho_- + \frac{\pi}{8} \right) + \frac{\pi Q}{4} \right\} + \cos \left\{ L \left(\rho_+ + \frac{3\pi}{8} \right) + \frac{3\pi Q}{4} \right\} \\
&= 0 \quad (22)
\end{aligned}$$

where $\rho_\pm = \arctan \{ (\sqrt{2} \pm 1) \tan \beta \}$.

For $L \gg 1$ and $\cos(2\beta)$ not too close to zero, the second term of (22) dominates and the solutions come where

$$\cos \left\{ L \left(\rho_+ + \frac{3\pi}{8} \right) + \frac{3\pi Q}{4} \right\} \approx 0.$$

After some rearrangement this leads to

$$\tan \beta_k \approx -\tan \frac{\pi}{4} \tan \frac{\pi}{8} \cot \left(K + \frac{\pi}{8} \right) \quad \text{with} \quad K = \frac{(4k + Q - 2)\pi}{4L}, \quad (23)$$

$k = 1, 2, \dots, [(3L - Q)/4]$ and e.g. $-\frac{\pi}{4} < \beta_k < \frac{\pi}{4}$. In contrast to the Z_3 -case, where in the approximation (11) the zeros were equally spaced in β , here this is

no longer the case: now the β_k are more dense in the center region. In the center $\beta \approx 0$ the oscillations of the ρ_+ -term in (22) are $(\sqrt{2}+1)/(\sqrt{2}-1) \approx 5.83$ times faster than those of the ρ_- -term. (20) and (23) are equivalent to

$$\cos \theta_k \equiv \frac{1+t_k^4}{1-t_k^4} \approx -\frac{\sin^4(K+\frac{\pi}{4})-\sin^4(K)}{\sin^4(K+\frac{\pi}{4})+\sin^4(K)}.$$

As in the Z_3 -case, in the thermodynamic limit, this approximation leads back to Baxter's ground-state energy formula.

5.2 The case Z_5 :

Transforming $t = \sin(\beta - \frac{\pi}{5})/\sin(\beta + \frac{\pi}{5})$, the condition

$$\mathcal{P}_Q \sim \mathcal{J}_0^L + \omega^{-Q} \mathcal{J}_1^L + \omega^Q \mathcal{J}_4^L + \omega^{-2Q} \mathcal{J}_2^L + \omega^{2Q} \mathcal{J}_3^L = 0$$

with $\mathcal{J}_k = t^4 + \omega^k t^3 + \omega^{2k} t^2 + \omega^{3k} t + \omega^{4k}$ takes the form

$$\begin{aligned} \frac{1}{2} \left(\frac{\cos \beta}{\cos \frac{\pi}{5}} \right)^{-L} + \cos \left\{ L \left(\beta + \frac{\pi}{5} \right) + \frac{2\pi Q}{5} \right\} \\ + B^{-L/2} \cos \left\{ L \left(\chi + \frac{2\pi}{5} \right) + \frac{4\pi Q}{5} \right\} = 0. \end{aligned} \quad (24)$$

where $\tan \chi = \cot \frac{\pi}{5} \cot \frac{\pi}{10} \tan \beta$ and $B = \frac{4 \sin^2 \beta + \sqrt{5} - 2}{2 \cos \frac{\pi}{5}}$.

The solutions come for $-\infty < s < 0$ corresponding to e.g. $-\frac{\pi}{5} < \beta < \frac{\pi}{5}$. Since for $L \gg 1$ and $|\beta|$ not too close to $\frac{\pi}{5}$ we have $1 \ll (\cos \beta / \cos \frac{\pi}{5})^{-L} \ll B^{-L/2}$, the solutions come close to where the cosine containing χ vanishes, i.e. at

$$\tan \beta_k \approx -\tan \frac{\pi}{5} \tan \frac{\pi}{10} \cot \left(K + \frac{\pi}{10} \right); \quad K = \frac{(10k + 2Q - 5)\pi}{10L}$$

and integers $k = 1, 2, \dots$ such that e.g. $|\beta_k| < \frac{\pi}{5}$. Equivalently, we can write

$$\cos \theta_k \equiv \frac{1+t_k^5}{1-t_k^5} \approx -\frac{\sin^5(K+\frac{\pi}{5})-\sin^5 K}{\sin^5(K+\frac{\pi}{5})+\sin^5 K}. \quad (25)$$

The Z_3 -formula (10) has the same form as the first terms of (24) since $2 \cos \beta = \cos \beta / \cos \frac{\pi}{3}$. The expressions for general Z_N will be given elsewhere.

5.3 The case Z_6 :

We conclude with the case Z_6 : As in the case of Z_4 , because $N = 6$ is even, we include a phase and transform $-\infty < s < 0$ to $-\frac{\pi}{6} < \beta < \frac{\pi}{6}$ using

$$s = t^6; \quad t = e^{i\pi/6} \frac{\sin(\beta - \frac{\pi}{6})}{\sin(\beta + \frac{\pi}{6})}.$$

Now $\mathcal{P}_Q(s) = 0$ (we don't give the explicit Z_6 formula for $\mathcal{P}_Q$, since this is easily generalized from the Z_5 -case) after some algebra leads to

$$\begin{aligned} \left(\frac{1-\alpha}{1+\alpha}\right)^{L/2} \cos\left\{L\left(\rho_- + \frac{\pi}{12}\right) + \frac{\pi Q}{6}\right\} + (1-\alpha)^{L/2} \cos\left\{L\left(\rho_0 + \frac{\pi}{4}\right) + \frac{\pi Q}{2}\right\} \\ + \cos\left\{L\left(\rho_+ + \frac{5\pi}{12}\right) + \frac{5\pi Q}{6}\right\} = 0 \end{aligned} \quad (26)$$

$$\text{with} \quad \alpha = \frac{\sqrt{3}}{2} \frac{\tan^2 \frac{\pi}{6} - \tan^2 \beta}{\tan^2 \frac{\pi}{6} + \tan^2 \beta} \quad \text{and}$$

$$\rho_0 = \arctan(\sqrt{3} \tan \beta); \quad \rho_{\pm} = \arctan\left\{\left(2 \pm \sqrt{3}\right) \sqrt{3} \tan \beta\right\}. \quad (27)$$

For $L \gg 1$ and α not too close to zero, the last term containing $L\rho_+$ dominates and the solutions β_k approach

$$\tan \beta_k \approx -\tan \frac{\pi}{6} \tan \frac{\pi}{12} \cot\left(K + \frac{\pi}{12}\right); \quad K = \frac{(6k + Q - 3)\pi}{6L}$$

and $k = 1, 2, \dots, m_E$ with $m_E = [(5L - Q)/6]$. We can rewrite this like (25).

6 Conclusions

In the analytic solution of the superintegrable chiral Z_N -Potts quantum chain the energy eigenvalues are determined by the zeros of special polynomials. These polynomials also play a central role in the representation theory of the Onsager algebra. In the thermodynamic limit, sums over functions of the zeros can be evaluated by contour integrations. For finite systems we can try to calculate the zeros numerically. However, applying standard computer routines directly to the explicit form of the polynomials will run into trouble beyond some chain length L , because the largest coefficients of the polynomials are growing exponentially with increasing L . So we were looking for an alternative and possibly analytic approach.

By a suitable change of variable we transform the equations for the zeros into trigonometric equations, for which approximate analytic solutions can be obtained. We give explicit formulae for the simplest polynomials.

In the case of Z_3 the solutions corresponding to the zeros of the polynomials are almost equidistant in the variable β . However, for Z_4 and higher, the density of zeros in β varies appreciably over the finite interval. Our formula describes the solutions, except those near at the boundaries, with exponential precision in L . The leading finite-size effects are due to the solutions at the extreme ends of the interval in β . For the case Z_3 we derive a special formula which describes also the extreme solutions, thus allowing to calculate finite-size energy eigenvalues for the low-lying sectors.

Acknowledgements

I am grateful to Roger E. Behrend for many fruitful discussions. Part of this work was performed during a visit of the author to SISSA, Trieste where it was supported by the TIMR-Network Contract FMRX-CT96-0012 of the European Commission. I thank G. Mussardo for his kind hospitality at SISSA.

References

- Albertini G., McCoy B.M. and Perk J.H.H. (1989): Phys.Lett. **A135** 159; Advanced Studies in Pure Math., Vol.19 (Kinokuniya-Academic) 1
 Albertini G. and McCoy B.M. (1990): Nucl.Phys. **B350** 745
 Au-Yang H., McCoy B.M., Perk J.H.H., Tang S. and Yan M.L. (1987): Phys.Lett. **A123** 219
 Baxter R.J. (1982): Exactly Solved Models in Statistical Mechanics, Academic Press
 Baxter R.J. (1988): Phys.Lett. **A133** 185
 Baxter R.J., Bazhanov V.V. and Perk J.H.H. (1990): Int.J.Mod.Phys. **B4** 803
 Baxter R.J. (1993): J.Stat.Phys. **73** 461
 Baxter R.J. (1994): J.Phys.A: Math.Gen. **27** 1837
 Baxter R.J. (1998): cond-mat/9808120
 Bazhanov V.V. and Stroganov Yu.G. (1990): J.Stat.Phys. **59** 799
 Dasmahapatra S., Kadem R. and McCoy B.M. (1993), Nucl.Phys. **B396** 506
 Davies B. (1990): J.Phys.A: Math.Gen. **23** 2245
 Dolan L. and Grady M. (1982): Phys. Rev. **D 25** 1587
 Gehlen G.v. and Rittenberg V. (1985): Nucl.Phys. **B257** 351
 Hamer C.J., Quispel G.R.W. and Batchelor M.T. (1987): J.Phys.A, Math.Gen. **20** 5677
 Onsager L. (1944): Phys. Rev. **65** 117
 O'Rourke M.J. and Baxter R.J. (1996): J.Stat.Phys. **82** 1
 Roan S.-S. (1991): preprint Max-Planck-Inst. für Math. Bonn MPI/91-70
 Woynarovich F. and Eckle H.-P. (1987): J.Phys.A, Math.Gen. **20** L97
 Zhou Y.-K. (1998): Nucl.Phys. **B522** 550